

Referências Bibliográficas

- [1] **Blender**. www.blender.com.br.
- [2] AXLER, S.; BOURDON, P. ; RAMEY, W. **Harmonic Function Theory**. Springer, 2001.
- [3] FLOATER, M. S. **Mean value coordinates**. Computer Aided Geometric Design, 20(1):19–27, 2003.
- [4] FLOATER, M. S.; KOS, G. ; REIMERS, M. **Mean value coordinates in 3d**. Computer Aided Geometric Design, 22:623–631, 2005.
- [5] JOSHI, P.; MEYER, M.; DEROSE, T.; GREEN, B. ; SANOCKI, T. **Harmonic coordinates for character articulation**. ACM Transactions on Graphics, 26(3), 2007. Artigo 71.
- [6] JU, T.; LIEPA, P. ; WARREN, J. **A general geometric construction of coordinates in a convex simplicial polytope**. Computer Aided Geometric Design, 24(3):161–178, 2007.
- [7] JU, T.; SCHAEFER, S. ; WARREN, J. **Mean value coordinates for closed triangular meshes**. ACM Transactions on Graphics, 24(3):561–566, 2005.
- [8] LAGE, M. **Estruturas de dados topológicos escalonáveis para variedades de dimensão 2 e 3**. Dissertação de Mestrado, Departamento de Matemática, PUC-Rio, <http://www.matmidia.mat.puc-rio.br/mlage/>, 2006.
- [9] LIPMAN, Y.; KOPF, J.; COHEN-OR, D. ; LEVIN, D. **Gpu-assisted positive mean value coordinates for mesh deformation**. Symposium on Geometry Processing, p. 117–123, 2007.
- [10] LIPMAN, Y.; LEVIN, D. ; COHEN-OR, D. **Green coordinates**. ACM Transactions on Graphics, 27(3), 2008. 78.
- [11] PINKALL, U.; POLTHIER, K. **Computing discrete minimal surfaces and their conjugates**. Experimental Mathematics, 1993.

- [12] RUSTAMOV, R. M.; LIPMAN, Y. ; FUNKHOUSER, T. **Interior distance using barycentric coordinates**. Symposium on Geometry Processing, 28(5), 2009.